

Östrogene in der Menopause

C. Jaurisch-Hancke
aus BDI aktuell 6/2007

- 1 Aas AM et al. **Blood glucose lowering by means of lifestyle intervention has different effects on adipokines as compared with insulin treatment in subjects with type 2 diabetes.** Diabetologia 2006; 49:872-80
- 2 Boule NG et al. **Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: a meta-analysis of controlled clinical trials,** JAMA 2001; 286:1218-27
- 3 Chiasson et al, **STOP-NIDDM Trial Research.** Lancet 359 [2002], No. 9323: 2072-2077
- 4 Dansinger, ML, et al., JAMA 2005; 293: 43-53
- 5 Deutsche Adipositas Gesellschaft, **Leitlinie zur Therapie der Adipositas**, Version 2006; www.adipositas-gesellschaft.de
- 6 **Diabetes Prevention Program - DPP**, NEJM, Volume 346; 2002, 393-403
- 7 Dormandy JA et al.: **Secondary prevention of macrovascular events in patients with type 2 diabetes in the Proactive study: a randomised controlled trial.** Lancet 2005; 366: 1279-89
- 8 The DREAM Trial Investigators, Lancet 2006; 368: 1096-1105
- 9 Goodyear, Lauri J Kahn, Barbara B: **Exercise, Glukosetransport und Insulinsensitivity**, Annu. Rev. Ned. 1998, 49: 235-61
- 10 Hu, FB. et al: **Diet, Lifestyle, and the risk of Type 2 Diabetes Mellitus in Women**, NEJM, 345: 790-797, 2001
- 11 Hu, FB et al., **Walking compared with vigorous physical activity and risk of type 2 diabetes in women: A prospective Study.** JAMA 1999; 282: 1433-9
- 12 Kahn, Steven E. et al.: **Glycemic durability of Rosiglitazon, Metformin or Glyburide monotherapy**, NEJM 350: 2427-2443, 2006
- 13 Klein, S. et al., **Weight management through lifestyle modification for the prevention and management of type 2 diabetes**, American Journal of Clinical Nutrition, Vol. 80, No. 2, 257-263, 2004
- 14 Lindström J et al, **Sustained reduction in the incidence of typ 2 diabetes by lifestyle intervention; follow up of the Finnish Diabetes Prevention Study**, Lancet 2006, 368:1673-9
- 15 Sigal, RJ et al: **Physical activity/exercise and Type 2 Diabetes.** Diabetes Care 2004; 27:2518-39
- 16 Toeller, M., **Evidenzbasierte Ernährungsempfehlungen**, Diabetes und Stoffwechsel, 14/2005
- 17 Tuomilehto, J., et al: NEJM 344 (2001), 1343-1350
- 18 UKPDS Group: **Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34).** Lancet 352 (1998) 854-865
- 19 UKPDS Group: **Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33).** Lancet 352 (1998) 837-853