

Primärdiagnostik der Herzinsuffizienz in Klinik und Praxis

S. Störk, S. Frantz, J. Bauersachs, G. Ertl, C. E. Angermann

aus BDI aktuell 10/2008

Literatur

- 1 Recommendations for exercise testing in chronic heart failure patients. *Eur Heart J* 2001; 22: 37-45
- 2 Cleland JG, Freemantle N, Ball SG. et al. The heart failure revascularisation trial (HEART). *Eur J Heart Fail* 2003; 5: 295-303
- 3 Cooper LT, Baughman KL, Feldman AM. et al. The role of endomyocardial biopsy in the management of cardiovascular disease. *Eur Heart J* 2007; 28: 3076-3093
- 4 Gillespie ND, McNeill G, Pringle T. et al. Cross sectional study of contribution of clinical assessment and simple cardiac investigations to diagnosis of left ventricular systolic dysfunction in patients admitted with acute dyspnoea. *BMJ* 1997; 314: 936-940
- 5 Hoppe UC, Bohm M, Dietz R. et al. [Guidelines for therapy of chronic heart failure]. *Z Kardiol* 2005; 94: 488-509
- 6 Januzzi JL, van Kimmenade R, Lainchbury J. et al. NT-proBNP testing for diagnosis and short-term prognosis in acute destabilized heart failure. *Eur Heart J* 2006; 27: 330-337
- 7 Latini R, Masson S, Anand IS. et al. Prognostic value of very low plasma concentrations of troponin T in patients with stable chronic heart failure. *Circulation* 2007; 116: 1242-1249
- 8 Maisel AS, Krishnaswamy P, Nowak RM. et al. Rapid measurement of B-type natriuretic peptide in the emergency diagnosis of heart failure. *N Engl J Med* 2002; 347: 161-167
- 9 Maisel AS, McCord J, Nowak RM. et al. Bedside B-Type natriuretic peptide in the emergency diagnosis of heart failure with reduced or preserved ejection fraction. *J Am Coll Cardiol* 2003; 41: 2010-2017
- 10 McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure. *N Engl J Med* 1971; 285: 1441-1446
- 11 Ommen SR, Nishimura RA, Appleton CP. et al. Clinical utility of Doppler echocardiography and tissue Doppler imaging in the estimation of left ventricular filling pressures.. *Circulation* 2000; 102: 1788-1794
- 12 Paulus WJ, Tschope C, Sanderson JE. et al. How to diagnose diastolic heart failure. *Eur Heart J* 2007; 28: 2539-2550
- 13 Redfield MM, Rodeheffer RJ, Jacobsen SJ. et al. Plasma brain natriuretic peptide to detect preclinical ventricular systolic or diastolic dysfunction. *Circulation* 2004; 109: 3176-3181
- 14 Ruskoaho H. Cardiac hormones as diagnostic tools in heart failure. *Endocr Rev* 2003; 24: 341-356
- 15 Senni M, Tribouilloy CM, Rodeheffer RJ. et al. Congestive heart failure in the community. *Circulation* 1998; 98: 2282-2289
- 16 Swedberg K, Cleland J, Dargie H. et al. Guidelines for the diagnosis and treatment of chronic heart failure: executive summary (update 2005). *Eur Heart J* 2005; 26: 1115-1140
- 17 Wheeldon NM, MacDonald TM, Flucker CJ. et al. Echocardiography in chronic heart failure in the community. *Q J Med* 1993; 86: 17-23
- 18 Willenheimer RB, Israelsson BA, Cline CM, Erhardt LR. Simplified echocardiography in the diagnosis of heart failure. *Scand Cardiovasc J* 1997; 31: 9-16
- 19 Wright SP, Doughty RN, Pearl A. et al. Plasma amino-terminal pro-brain natriuretic peptide and accuracy of heart-failure diagnosis in primary care. *J Am Coll Cardiol* 2003; 42: 1793-1800
- 20 Wu AH, Smith A, Wiecezorek S. et al. Biological variation for N-terminal pro- and B-type natriuretic peptides and implications for therapeutic monitoring of patients with congestive heart failure. *Am J Cardiol* 2003; 92: 628-631
- 21 Zaphiriou A, Robb S, Murray-Thomas T. et al. The diagnostic accuracy of plasma BNP and NTproBNP in patients referred from primary care with suspected heart failure. *Eur J Heart Fail* 2005; 7: 537-541