

Gewichtsreduktion bei Adipositas: Welche Rolle spielen die Gene?

C. Holzapfel, H. Hauner

aus BDI aktuell 7/2009

Literatur

- 1 Alhassan S, Kim S, Bersamin A, King AC, Gardner CD. Dietary adherence and weight loss success among overweight women: results from the A to Z weight loss study. *Int J Obes* 2008; 32: 985–991
- 2 Arner P. Genetic variance and lipolysis regulation: implications for obesity. *Ann Med* 2001; 33: 542–546
- 3 Benecke H, Topak H, von zur Mühlen A, Schuppert F. A study on the genetics of obesity: influence of polymorphisms of the beta-3-adrenergic receptor and insulin receptor substrate 1 in relation to weight loss, waist to hip ratio and frequencies of common cardiovascular risk factors. *Exp Clin Endocrinol Diabetes* 2000; 108: 86–92
- 4 Bouchard C, Tremblay A. Genetic influences on the response of body fat and fat distribution to positive and negative energy balances in human identical twins. *J Nutr* 1997; 127: 943S–947S
- 5 Bouchard C, Tremblay A, Després JP et al. The response to exercise with constant energy intake in identical twins. *Obes Res* 1994; 2: 400–410
- 6 Clement K, Vaisse C, Lahlou N et al. A mutation in the human leptin receptor gene causes obesity and pituitary dysfunction. *Nature* 1998; 392: 398–401
- 7 Corella D, Qi L, Sorli JV et al. Obese subjects carrying the 11482G>A polymorphism at the perilipin locus are resistant to weight loss after dietary energy restriction. *J Clin Endocrinol Metab* 2005; 90: 5121–5126
- 8 Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA* 2005; 293: 43–53
- 9 Dina C, Meyre D, Gallina S et al. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet* 2007; 39: 724–726
- 10 Farooqi IS, Matarese G, Lord GM et al. Beneficial effects of leptin on obesity, T cell hyporesponsiveness, and neuroendocrine/metabolic dysfunction of human congenital leptin deficiency. *J Clin Invest* 2002; 110: 1093–1103
- 11 Frayling TM, Timpson NJ, Weedon MN et al. A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 2007; 316: 889–894
- 12 Frey UH, Hauner H, Jöckel KH, Manthey I, Brockmeyer N, Siffert W. A novel promoter polymorphism in the human gene GNAS affects binding of transcription factor upstream stimulatory factor 1, Gas protein expression and body weight regulation. *Pharmacogenet Genomics* 2008; 18: 141–151
- 13 Fumeron F, Durack-Bown I, Betoulle D et al. Polymorphisms of uncoupling protein (UCP) and beta 3 adrenoreceptor genes in obese people submitted to a low calorie diet. *Int J Obes Relat Metab Disord* 1996; 20: 1051–1054
- 14 Gerken T, Girard CA, Tung YC et al. The obesity-associated FTO gene encodes a 2-oxoglutarate-dependent nucleic acid demethylase. *Science* 2007; 318: 1469–1472
- 15 Hainer V, Stunkard AJ, Kunesová M, Parízková J, Stich V, Allison DB. A twin study of weight loss and metabolic efficiency. *Int J Obes Relat Metab Disord* 2001; 25: 533–537
- 16 Hainer V, Zamrazilová J, Spálová J et al. Role of hereditary factors in weight loss and its maintenance. *Physiol Res* 2008; 57: S1–S15
- 17 Harper ME, Dent R, Monemdjou S et al. Decreased mitochondrial proton leak and reduced expression of uncoupling protein 3 in skeletal muscle of obese diet-resistant women. *Diabetes* 2002; 51: 2459–2466

- 18** Hauner H, Meier M, Jöckel KH, Frey UH, Siffert W. Prediction of successful weight reduction under sibutramine therapy through genotyping of the G-protein $\beta 3$ subunit gene (GNB3) C825T polymorphism. *Pharmacogenetics* 2003; 13: 453–459
- 19** Hauner H, Röhrig K, Siffert W. Effects of the G-Protein $\beta 3$ subunit 825T allele on adipogenesis and lipolysis in cultured human preadipocytes and adipocytes. *Horm Metab Res* 2002; 34: 475–480
- 20** Hauner H, Buchholz G, Hamann A et al. Prävention und Therapie der Adipositas. Evidenzbasierte Leitlinie der Deutschen Adipositas-Gesellschaft, der Deutschen Diabetes-Gesellschaft, der Deutschen Gesellschaft für Ernährung, der Deutschen Gesellschaft für Ernährungsmedizin. 2007 www.adipositas-gesellschaft.de
- 21** Haupt A, Thamer C, Machann J et al. Impact of variation in the FTO gene on whole body fat distribution, ectopic fat, and weight loss. *Obesity* 2008; 16: 1969–1972
- 22** Heitmann BL, Lissner L, Sorensen TIA, Bengtsson C. Dietary fat intake and weight gain in women genetically predisposed for obesity. *Am J Clin Nutr* 1995; 61: 1213–1217
- 23** Heshka S, Anderson JW, Atkinson RL et al. Weight loss with self-help compared with a structured commercial program: a randomized trial. *JAMA* 2003; 289: 1792–1798
- 24** Kawaguchi H, Masuo K, Katsuya T et al. Beta2- and beta3-adrenoceptor polymorphisms relate to subsequent weight gain and blood pressure elevation in obese normotensive individuals. *Hypertens Res* 2006; 29: 951–959
- 25** Lindi V, Sivenius K, Niskanen L, Laakso M, Uusitupa MI. Effect of the Pro12Ala polymorphism of the PPAR-gamma2 gene on long-term weight change in Finnish non-diabetic subjects. *Diabetologia* 2001; 44: 925–926
- 26** Mammès O, Betoulle D, Aubert R et al. Novel polymorphisms in the 5' region of the LEP gene. Association with leptin levels and response to low-calorie diet in human obesity. *Diabetes* 1998; 47: 487–489
- 27** Mammès O, Aubert R, Betoulle D et al. LEPR gene polymorphisms: associations with overweight, fat mass and response to diet in women. *Eur J Clin Invest* 2001; 31: 398–404
- 28** Moreno-Aliaga MJ, Santos JL, Marti A, Martinez JA. Does weight loss prognosis depend on genetic make-up? *Obes Rev* 2005; 6: 155–168
- 29** Müller TD, Hinney A, Scherag A et al. Fat mass and obesity associated gene (FTO): no significant association of variant rs9939609 with weight loss in a lifestyle intervention and lipid metabolism markers in German obese children and adolescents. *BMC Med Genet* 2008; 9: 85
- 30** Näslund E, Andersson I, Degerblad M et al. Associations of leptin, insulin resistance and thyroid function with long-term weight loss in dieting men. *J Intern Med* 2000; 248: 299–308
- 31** Nicklas BJ, van Rossum EFC, Berman DM, Ryan AS, Dennis KE, Shuldiner AR. Genetic variation in the peroxisome proliferator-activated receptor-g2 gene (Pro12Ala) affects metabolic responses to weight loss and subsequent weight regain. *Diabetes* 2001; 50: 2172–2176
- 32** Peters WR, Macmurry JP, Walker J, Giese RJJr, Comings DE. Phenylethanolamine N-Methyltransferase G-148A genetic variant and weight loss in obese women. *Obes Res* 2003; 11: 415–419
- 33** Pooley EC, Fairburn CG, Cooper Z, Sodhi MS, Cowen PJ, Harrison PJ. A 5-HT2C receptor promoter polymorphism (HTR2C-759C/T) is associated with obesity in women, and with resistance to weight loss in heterozygotes. *Am J Med Genet* 2004; 126B: 124–127
- 34** Rawson ES, Nolan A, Silver K, Shuldiner AR, Poehlman ET. No effect of the Trp64Arg beta(3)-adrenoreceptor gene variant on weight loss, body composition, or energy expenditure in obese, caucasian postmenopausal women. *Metabolism* 2002; 51: 801–805
- 35** Reinehr T, Hinney A, Nguyen TT, Hebebrand J. Evidence of an influence of a polymorphism near the INSIG2 on weight loss during a lifestyle intervention in obese children and adolescents. *Diabetes* 2008; 57: 623–626
- 36** Reinehr T, Hebebrand J, Friedel S et al. Lifestyle intervention in obese children with variations in the melanocortin 4 receptor gene. *Obesity* 2009; 17: 382–389
- 37** Rucker D, Padwal R, Li SK, Curioni C, Lau DCW. Long-term pharmacotherapy for obesity and overweight: updated meta-analysis. *BMJ* 2007; 335: 1194–1199 (Erratum in *BMJ* 2007;335)

- 38** Sakane N, Yoshida T, Umekawa T, Kogure A, Takakura Y, Kondo M. Effects of Trp64Arg mutation in the beta-3 adrenergic receptor gene on weight loss, body fat distribution, glycemic control, and insulin resistance in obese type 2 diabetic patients. *Diabetes Care* 1997; 20: 1887–1890
- 39** Scuteri A, Sanna S, Chen WM et al. Genome-wide association scan shows genetic variants in the FTO gene are associated with obesity-related traits. *PLOS Genet* 2007; 3: e115
- 40** Shai I, Schwarzfuchs D, Henkin Y et al. Weight loss with a low-carbohydrate, mediterranean, or low-fat diet. *N Engl J Med* 2008; 359: 229–241
- 41** Shin HD, Kim KS, Cha MH, Yoon Y. The effects of UCP-1 polymorphisms on obesity phenotypes among Korean female subjects. *Biochem Biophys Res Commun* 2005; 335: 624–630
- 42** Shiwaku K, Nogi A, Anuurad E et al. Difficulty in losing weight by behavioral intervention for women with Trp64Arg polymorphism of the beta-3 adrenergic receptor gene. *Int J Obes Relat Metab Disord* 2003; 27: 1028–1036
- 43** Sorensen TI, Boutin P, Taylor MA et al. Genetic polymorphisms and weight loss in obesity: a randomised trial of hypo-energetic high- versus low-fat diets. *Plos Clin Trials* 2006; 1: e12
- 44** Stunkard AJ, Foch TT, Hrubec Z. A twin study of human obesity. *JAMA* 1986a; 256: 51–54
- 45** Stunkard AJ, Sorensen TI, Hanis C et al. An adoption study of human obesity. *N Engl J Med* 1986b; 314: 193–198
- 46** Svetkey LP, Stevens VJ, Brantley PJ et al. Comparison of strategies for sustaining weight loss: the weight loss maintenance randomized controlled trial. *JAMA* 2008; 299: 1139–1148
- 47** Tchernof A, Starling RD, Turner A et al. Impaired capacity to lose visceral adipose tissue during weight reduction in obese postmenopausal women with the Trp64Arg beta3-adrenoceptor gene variant. *Diabetes* 2000; 49: 1709–1713
- 48** Thorleifsson G, Walters GB, Gudbjartsson DF et al. Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat Genet* 2009; 41: 18–24
- 49** Verdich C, Toubro S, Buemann B et al. Leptin levels are associated with fat oxidation and dietary-induced weight loss in obesity. *Obes Res* 2001; 9: 452–461
- 50** Vogels N, Mariman ECM, Bouwman FG, Kester ADM, Diepvens K, Westerterp-Plantenga MS. Relation of weight maintenance and dietary restraint to peroxisome proliferator-activated receptor g2, glucocorticoid receptor, and ciliary neurotrophic factor polymorphisms. *Am J Clin Nutr* 2005; 82: 740–746
- 51** Westberg L, Bah J, Rastam M et al. Association between a polymorphism of the 5-HT2C receptor and weight loss in teenage girls. *Neuropsychopharmacology* 2002; 26: 789–793
- 52** Willer CJ, Speliotes EK, Loos RJF et al. Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nat Genet* 2009; 41: 25–34
- 53** Yoon Y, Park BL, Cha MH et al. Effects of genetic polymorphisms of UCP2 and UCP3 on very low calorie diet-induced body fat reduction in Korean female subjects. *Biochem Biophys Res Commun* 2007; 359: 451–456
- 54** Farooqi JS, O'Rahilly S et al. Genetics of obesity in humans. *Endocrine Reviews* 2006; 27: 710–718
- 55** Loos RJF, Lindgren CM, Li S et al. Common variants near MC4R are associated with fat mass, weight and risk of obesity. *Nat Genet* 2008; 40: 768–775